

Supplementary Table 2. Density of premonitory urges (PUs) in all 95 body regions investigated

95 body regions	<i>n</i> * (%†)
R_Ant_Forehead	448 (4.5)
L_Ant_Forehead	445 (4.5)
R_Ant_Cheek	417 (4.2)
L_Ant_Cheek	401 (4)
C_Ant_Mouth	385 (3.9)
L_Ant_Chest	228 (2.3)
R_Ant_Chest	228 (2.3)
L_Ant_Throat	226 (2.3)
R_Ant_Abdomen	222 (2.2)
L_Ant_Abdomen	221 (2.2)
C_Ant_Throat	216 (2.2)
R_Ant_Throat	206 (2.1)
R_Ant_Eye	199 (2)
C_Ant_Nose	189 (1.9)
L_Ant_Eye	186 (1.9)
R_Ant_Hand	179 (1.8)
R_Post_Occiput	168 (1.7)
L_Ant_Hand	162 (1.6)
L_Post_Occiput	162 (1.6)
C_Ant_Chest	161 (1.6)
L_Ant_Shoulder	160 (1.6)
L_Post_Shoulder	159 (1.6)
C_Ant_Forehead	158 (1.6)
L_Post_Neck	158 (1.6)
R_Ant_Shoulder	157 (1.6)
R_Post_Shoulder	152 (1.5)
R_Post_Neck	148 (1.5)
C_Ant_Chin	124 (1.3)
C_Post_Occiput	112 (1.1)
R_Ant_Foot	111 (1.1)
R_Ant_Chin	110 (1.1)
R_Ant_Upperarm	109 (1.1)
L_Ant_Foot	108 (1.1)
C_Post_Neck	102 (1)
L_Ant_Upperarm	102 (1)
C_Ant_Abdomen	96 (1)
L_Ant_Chin	95 (1)
L_Ant_LowerForearm	91 (0.9)
R_Post_Hand	90 (0.9)
L_Post_Hand	88 (0.9)
C_Ant_Upperlip	86 (0.9)
R_Ant_LowerForearm	84 (0.8)
L_Ant_Wrist	75 (0.8)
R_Post_Foot	73 (0.7)
L_Post_Foot	72 (0.7)
R_Ant_Ankle	72 (0.7)
R_Post_Abdomen	71 (0.7)
L_Ant_Ankle	69 (0.7)
L_Post_Upperarm	69 (0.7)
L_Post_Abdomen	68 (0.7)
R_Ant_Wrist	67 (0.7)
R_Post_Upperarm	67 (0.7)
L_Ant_Elbow	66 (0.7)
R_Ant_Pelvis	65 (0.7)
L_Ant_Pelvis	61 (0.6)
L_Post_UpperBack	61 (0.6)
R_Post_UpperBack	60 (0.6)
L_Post_Pelvis	54 (0.5)
L_Post_Forearm	50 (0.5)
L_Post_Wrist	50 (0.5)
R_Ant_Elbow	50 (0.5)
R_Ant_Ear	48 (0.5)
R_Post_Forearm	48 (0.5)
R_Post_Wrist	48 (0.5)
L_Ant_Ear	47 (0.5)
R_Post_Pelvis	47 (0.5)
R_Ant_Knee	43 (0.4)
R_Post_Ankle	43 (0.4)
R_Ant_Tibia	42 (0.4)
L_Ant_Tibia	41 (0.4)
L_Post_Ankle	41 (0.4)
L_Post_Knee	41 (0.4)
L_Ant_Knee	40 (0.4)
R_Post_Knee	39 (0.4)
L_Post_Elbow	37 (0.4)
L_Post_Tibia	35 (0.4)
R_Post_Tibia	33 (0.3)
L_Ant_UpperThigh	32 (0.3)
R_Ant_UpperForearm	32 (0.3)
R_Post_Elbow	32 (0.3)
R_Post_LowerBack	32 (0.3)
L_Post_LowerBack	30 (0.3)
L_Post_Thigh	27 (0.3)
R_Ant_UpperThigh	26 (0.3)
R_Post_Thigh	25 (0.3)
C_Post_Abdomen	19 (0.2)
R_Ant_LowerThigh	19 (0.2)
C_Post_UpperThorax	15 (0.2)
R_Post_Ear	14 (0.1)
C_Ant_Pelvis	13 (0.1)
L_Ant_LowerThigh	12 (0.1)
L_Post_Ear	12 (0.1)
C_Post_LowerThorax	10 (0.1)
C_Post_Pelvis	10 (0.1)
L_Ant_UpperForearm	8 (0.1)

*absolute numbers are based on the number of marks per body area of all tics and all participants experiencing PU (*n* = 232); †percentages are based on the number of marks per body area of all tics and all participants (*n* = 232) divided by the number of marks of all body areas of all tics and all participants (*n* = 232). C, center; L, left; R, right; Ant, anterior; Post, posterior.