

Supplementary Table 4. Changes of SQ-CPM during follow-up and evaluation according to the MDS-PD diagnostic criteria of the patient POD01 with DLB

Item	POD01		
F/U year	1	2	3
F/U SQ-CPM			
1. Clinical prodromal marker screening			
1-1. Hyposmia	N	N	Y
1-2. Constipation	N	N	Y
1-2-1. Frequency of defecation (day/frequency)	1.00	1.00	1.50
1-3. RBD1Q	N	N	N
2. Motor symptom screening			
2-1. Slowness	Y	Y	Y
2-2. Stooped posture	N	Y	Y
2-3. Rigidity	N	N	Y
2-4. Gait instability	N	N	Y
MDS-PD diagnostic criteria			
Step 1. Parkinsonism	N	N	Y
Bradykinesia		N	Y
Rigidity		N	Y
Rest tremor		N	N
Step 2. Supportive criteria		0	0
Step 3. Absolute exclusion criteria		0	0
Step 4. Red flags		0	3 (DLB)
Step 5. Diagnosis of PD		3	

DLB, dementia with Lewy bodies; PD, Parkinson's disease; SQ-CPM, screening questionnaire of clinical prodromal markers; RBD1Q, rapid eye movement sleep behavior disorder single-question screen; MDS, movement disorder society; POD, postoperative delirium; F/U, follow-up.