

Supplementary Table 1. Clinical scales to assess orthostatic dizziness

	Total (<i>n</i> = 42)	HCs (<i>n</i> = 11)	PD-OH(-) (<i>n</i> = 21)	PD-OH(+) (<i>n</i> = 10)	<i>p</i> value	HC vs. PD-OH(-)	HC vs. PD-OH(+)	PD-OH(-) vs. PD-OH(+)
SCOPA-AUT	7.0 ± 7.1	2.9 ± 3.6	9.5 ± 8.3	6.2 ± 5.1	0.029	0.007	0.099	0.492
SCOPA-AUT-CV ≥1	25 (59.5)	1 (9.1)	16 (76.2)	8 (80.0)	<0.001	<0.001	0.002	1.000
OH-SA (total)	5.1 ± 9.2	0.3 ± 0.6	8.1 ± 11.9	4.2 ± 4.4	0.021	0.012	0.013	0.724
OH-SA item 1 ≥1	21 (50.0)	2 (18.2)	12 (57.1)	7 (70.0)	0.063	0.061	0.030	0.697

Values are means ± standard deviation or *n* (%). Statistical analyses were performed using Kruskal–Wallis test or Fisher's exact test. Post hoc analyses were performed using Mann–Whitney test and chi-squared test.

HCs, healthy controls; PD-OH(-), Parkinson's disease patients without orthostatic hypotension; PD-OH(+), Parkinson's disease patients with orthostatic hypotension; SCOPA-AUT, Scale for Outcomes in Parkinson's Disease-Autonomic; SCOPA-AUT-CV, Scale for Outcomes in Parkinson's Disease-Autonomic-cardiovascular; OH-SA, Orthostatic Hypotension Symptom Assessment.